

STELLAR DYNAMICS AT THE GALACTIC CENTER WITH AN EXTREMELY LARGE TELESCOPE

NEVIN N. WEINBERG¹, MILOŠ MILOSAVLJEVIĆ¹, AND ANDREA M. GHEZ²*Draft version February 2, 2008*

ABSTRACT

We discuss physical experiments achievable via the monitoring of stellar dynamics near the massive black hole at the Galactic center with a diffraction-limited, next generation, extremely large telescope (ELT). Given the likely observational capabilities of an ELT and what is currently known about the stellar environment at the Galactic Center, we synthesize plausible samples of stellar orbits around the black hole. We use the Markov Chain Monte Carlo method to evaluate the constraints that the monitoring of these orbits will place on the matter content within the dynamical sphere of influence of the black hole. We express our results as functions of the number N of stars with detectable orbital motions and the astrometric precision $\delta\theta$ and spectroscopic precision δv at which the stellar proper motions and radial velocities are monitored. Our results are easily scaled to different telescope sizes and precisions. For $N = 100$, $\delta\theta = 0.5$ mas, and $\delta v = 10$ km s⁻¹—a conservative estimate of the capabilities of a 30 meter telescope—we find that if the extended matter distribution enclosed by the orbits at 0.01 pc has a mass greater than $\sim 10^3 M_\odot$, it will produce measurable deviations from Keplerian motion. Thus, if the concentration of dark matter at the Galactic Center matches theoretical predictions, its influence on the orbits will be detectable. We also estimate the constraints that will be placed on the mass of the black hole and on the distance to the Galactic Center, and find that both will be measured to better than $\sim 0.1\%$. We discuss the significance of knowing the distance to within a few parsecs and the importance of this parameter for understanding the structure of the Galaxy. We demonstrate that the lowest-order relativistic effects, such as the prograde precession, will be detectable if $\delta\theta \lesssim 0.5$ mas. Barring the favorable discovery of a star on a highly compact, eccentric orbit, the higher-order effects, including the frame dragging due to the spin of the black hole, will require $\delta\theta \lesssim 0.05$ mas. Finally, we calculate the rate at which monitored stars experience detectable nearby encounters with background stars. The encounters probe the mass function of stellar remnants that accumulate near the black hole. We find that ~ 30 such encounters will be detected over a ten year baseline for $\delta\theta = 0.5$ mas.

Subject headings: astrometry — black hole physics — Galaxy: center — Galaxy: kinematics and dynamics — infrared: stars

1. INTRODUCTION

Observational programs with ten meter class telescopes, including the W. M. Keck Observatory and the Very Large Telescope (VLT), have yielded a wealth of information on the stellar content inside the sphere of influence of the massive black hole at the Galactic center (GC; Ghez et al. 1998; Gezari et al. 2002; Hornstein et al. 2002; Figer et al. 2003; Genzel et al. 2003a; Ghez et al. 2003b; Schödel et al. 2003). The black hole is located at the center of a compact stellar cluster that has been the target of observational surveys for a decade (e.g., Krabbe et al. 1995; Figer et al. 2000; Gezari et al. 2002). Near-infrared monitoring with speckle and adaptive optics techniques has recently enabled complete orbital reconstruction of several stellar sources orbiting the black hole (Eckart et al. 2002; Schödel et al. 2002, 2003; Ghez et al. 2003b). Sources have been monitored with astrometric errors of a few milli-arcseconds (Ghez et al. 2003a; Schödel et al. 2003), and radial velocity errors < 50 km s⁻¹ (Eisenhauer et al. 2003; Ghez et al. 2003a), allowing the detection of the accelerated proper motions of ~ 10 stars. One of these stars has an orbital period of only ~ 15 yr (Ghez et al. 2003b; Schödel et al. 2003).

The presence of a dark mass at the center of the Galaxy could in principle be inferred from the static nature of the ra-

dio source Sgr A* located at the center of the stellar cluster (Backer & Sramek 1999; Reid et al. 1999). Nevertheless, it is the stars with the shortest orbital periods that have provided unequivocal proof of the existence of a massive black hole and a measurement of its mass of $\sim 4 \times 10^6 M_\odot$ (Ghez et al. 2003b; Schödel et al. 2003). Since, for a fixed angular scale, the orbital periods are proportional to $R_0^{3/2} M_{\text{bh}}^{-1/2}$ and the radial velocities are proportional to $R_0^{-1/2} M_{\text{bh}}^{1/2}$ where R_0 is the heliocentric distance to the black hole and M_{bh} is its mass, the two parameters are not degenerate and can be determined independently (Eisenhauer et al. 2003).

In spite of the quality of elementary data available about the black hole and the bright stellar sources, the matter content in the vicinity of the black hole remains unknown. The observed stellar sources may represent only a fraction of the total matter content. Since the radial diffusion time $\sim 10^{8-9}$ yr is shorter than the age of the bulge, a large number of massive compact remnants ($5-10 M_\odot$ black holes) could have segregated into, and may dominate the matter density inside the dynamical sphere of influence of the black hole (Morris 1993; Miralda-Escudé & Gould 2000). Furthermore, adiabatic growth of the massive black hole could have compressed a pre-existing distribution of cold dark matter (CDM) (Ipser & Sikivie 1987; Quinlan, Hernquist, & Sigurdsson 1995; Gondolo & Silk 1999) and stars (Peebles 1972; Young 1980) into a dense “spike”. A variety of dynamical processes, however, are capable of destroying such a spike (Ullio, Zhao, & Kamionkowski 2001; Merritt et al. 2002;

¹ Theoretical Astrophysics, California Institute of Technology, Pasadena, CA 91125; nnw, milos@tapir.caltech.edu

² Department of Physics and Astronomy, University of California, Los Angeles, CA 90095-1562; ghez@astro.ucla.edu

Gnedin & Primack 2003; Merritt 2003). A sustained CDM spike would have implications for the detection of annihilation radiation for the CDM models in which the CDM consists of weakly-interacting massive particles (WIMPs).

The most complete catalogue of stars in the central parsecs was compiled by Genzel et al. (2000) and Schödel et al. (2003). In a survey of the stellar sources, Genzel et al. (2003a) infer a spatial number density of $n(r) \propto r^{-1.4}$ over the radial range $0.004 < r < 0.4$ pc. Their sample was 50% complete for stars brighter than $K \sim 18$, where completeness is defined as the percentage of stars in the field of view that are detectable and thus included in the sample. Expressed in terms of stellar mass, the sample is 50% complete for masses $m \gtrsim 3M_\odot$, assuming stars on the main sequence, a distance to the GC of 8.0 kpc (Reid 1993) and K -band extinction of 3.3 mag (Rieke, Rieke, & Paul 1989). A picture is emerging in which the brightest stars in the Central Cluster (< 0.03 pc) are young, main-sequence stars with apparent magnitudes $K > 13$ and masses $10 - 15M_\odot$. The stars outside 0.03 pc appear to be spectroscopically and kinematically distinct. They span a larger range of magnitudes $K \gtrsim 10$ and contain ~ 40 mass-losing Wolf-Rayet stars (e.g., Genzel et al. 2003a and R. Genzel, private communication). Unlike the Central Cluster, these stars appear to belong to twin, misaligned stellar disks (Levin & Beloborodov 2003; Genzel et al. 2003a).

The formation of the observed young stars with $\times 100$ larger specific binding energies relative to the black hole than that of the nearest observed accumulation of molecular gas (e.g., Jackson et al. 1993) presents a challenge to star formation theories and is a persistent puzzle (e.g., Morris 1993; Ghez et al. 2003a; Genzel et al. 2003a). A number of mechanisms for the formation and migration of stars in the tidal field of the massive black hole have been proposed (Gerhard 2001; Gould & Quillen 2003; Hansen & Milosavljević 2003; Levin & Beloborodov 2003; Kim & Morris 2003; Milosavljević & Loeb 2004). While the mechanisms have important implications, they are also each deficient in at least one way.

There is a dearth of giants in the GC region (Eckart et al. 1995). Recently, Figer et al. (2003) measured the radial velocities of 85 cool, normal giant stars with projected distances from the central region between 0.1–1 pc. They find nearly complete deficiency of giants with large radial velocities ($V_{\text{rad}} > 200 \text{ km s}^{-1}$). Since a star in a circular orbit at a distance of 0.1 pc from the black hole has velocity $\sim 400 \text{ km s}^{-1}$, the absence of any such stars with comparable radial velocities indicates that the observed giants are indeed limited to the region outside the central ~ 0.5 pc.

While the measured stellar density profile of the Galactic bulge is consistent with that of a singular isothermal sphere (Becklin 1968), the profile in the central parsec is not well known, especially for the lower-mass stellar populations. Assuming relaxation that is driven by two-body processes, Bahcall & Wolf (1976) showed that the equilibrium phase space distribution for a population of equal mass stars is a power law in density $\rho \propto r^{-7/4}$. For a multimass distribution the lighter stars are less centrally concentrated, resulting in a power-law profile that ranges from $r^{-3/2}$ for the least massive species to $r^{-7/4}$ for the most (Bahcall & Wolf 1977; Murphy, Cohn, & Durisen 1991). A coeval family of stars in the central region has reached equilibrium only if it is older

than the relaxation time

$$t_E \sim \frac{\sigma^3}{G^2 m_* \rho \ln \Lambda} \\ \approx 2 \times 10^8 \text{ yr} \left(\frac{r}{1 \text{ pc}} \right)^{1/4} \left(\frac{M_{\text{bh}}}{4 \times 10^6 M_\odot} \right)^{3/2} \\ \times \left(\frac{m_*}{10 M_\odot} \right)^{-1} \left(\frac{\rho_{1\text{pc}}}{2 \times 10^5 M_\odot \text{ pc}^{-3}} \right)^{-1} \left(\frac{\ln \Lambda}{10} \right)^{-1} \quad (1)$$

where σ is the local linear stellar velocity dispersion, m_* is the mass of a typical field star, ρ is the local stellar density, and $\ln \Lambda$ is the Coulomb logarithm.

Since the main sequence lifetime of stars more massive than $\sim 2M_\odot$ is shorter than t_E , young massive stars in the GC are not relaxed; their distribution is primarily a reflection of their formative conditions. While lower mass dwarf stars are sufficiently old to be relaxed in the central potential, their distribution in the innermost region could be affected by an abundance of stellar mass black holes ($5 - 10M_\odot$). As products of normal stellar evolution, stellar mass black holes sink in the potential of the massive black hole (Morris 1993; Miralda-Escudé & Gould 2000) and displace the less massive stars and remnants.

Speckle imaging and more recently adaptive optics with the Keck and VLT have provided several milliarcsecond astrometry, enabling the detection of proper motions within the inner 0.5 pc and accelerated proper motions of ~ 10 stars within the inner 0.05 pc. Radial velocities with spectroscopic precisions of $\delta v \sim 30 \text{ km s}^{-1}$ have also been obtained for the star S0-2, which has been monitored for over 70% of its orbit including pericenter passage at ~ 130 AU from the black hole. These observations have enabled the black hole mass and GC distance to be measured to within $\sim 10\%$ (Ghez et al. 2003b; Schödel et al. 2003).

Here we examine the extent to which one can probe the GC potential by monitoring stars with a diffraction-limited, next generation, extremely large telescope (ELT). As compared with current 10 m class telescopes, the finer angular resolution of an ELT enables the orbital motions of many more stars to be detected, each at greater astrometric precision, $\delta\theta$, and spectroscopic precision, δv . Given the range of possible sizes of future telescope and given the uncertainties in the ultimate capabilities of a specific telescope class (e.g., 30 meter telescopes) we choose to express our results not as functions of the ELT aperture but rather as functions of $\delta\theta$, δv , and the number N of stars with detectable orbital motions. We take $\delta\theta = 0.5 \text{ mas}$ and $\delta v = 10 \text{ km s}^{-1}$ as our fiducial model, corresponding to a conservative estimate of the capabilities of a telescope with a $D = 30 \text{ m}$ aperture. We show that N scales with telescope aperture as $N \simeq 100(D/30 \text{ m})^2$. We demonstrate that with an ELT one can measure the density profile of a dark matter spike and those general relativistic effects that scale as $(v/c)^2$, where v is the speed of a star and c is the speed of light. Furthermore, we show that the distance to the GC will be measured to remarkable precision. This will help place tight constraints on models of the overall Galactic structure. We also show that with an ELT one can detect the gravitational interactions between monitored stars and the background massive stellar remnants that accumulate near the central black hole. Such interactions may probe the mass function of the stellar mass black holes thought to dominate the matter density in the region.

The paper is organized as follows. In § 2 we calculate the

number of stars with accelerated proper motions that can be monitored with a given ELT based on its astrometric, spectroscopic, and confusion limits. We also describe a realistic monitoring program and demonstrate that confusion with the infrared emission from Sgr A* is unlikely to affect an ELT's ability to measure stellar motions. In § 3 we model the orbital data and estimate the magnitude of various non-Keplerian effects including Newtonian retrograde precession due to extended matter, relativistic prograde precession, precession induced by the coupling of orbits to the spin of the black hole, and the Roemer time delay. In § 3.6 we consider the effect of stellar interactions on the motion of the monitored stars. Specifically, we estimate the rate at which discrete stellar encounters result in detectable changes of orbital motions. In § 4 we discuss a method for generating mock ELT orbital data and describe a computational technique for estimating uncertainties in the orbital parameters. The results of our calculations are given in § 5. Finally, in § 6 we discuss astrophysical applications of the proposed observations.

2. OBSERVING STARS IN THE CENTRAL ARCSECOND WITH AN ELT

The purpose of this section is to estimate the number and distribution of stars whose orbital motions can be detected with an ELT and to determine the astrometric and spectroscopic precision to which their motions can be measured (§ 2.1). The latter are determined by the specifications of the telescope and the properties of the stellar population at the GC. Several factors complicate the monitoring of orbits within the central arcsecond. The greatest obstacle to detecting and following hitherto unseen stars is stellar crowding. Light contamination from nearby bright stars as well as the light from underlying faint stars flood the pixel elements and impose a limit to the faintest detectable star. In § 2.2 we estimate the minimum luminosity permitted by the crowding and thereby obtain an estimate of the number count of stars with observable orbital motions.

2.1. Astrometric and Spectroscopic Limit

With adaptive optics, an ELT will operate near its diffraction limit in the K -band. By determining the centroid of images, the measured relative positions of stars are a factor of ~ 20 – 40 more definite than the images' diffraction limit. For instance, the diffraction limit of Keck is ~ 50 mas while the astrometric error of a bright star near the GC as seen by Keck is ~ 1 – 2 mas. Naively, the expected astrometric limit of an ELT with $D = 30$ m is therefore $\delta\theta_{30} \sim 0.5$ mas.

In practice, the astrometric limit achievable with adaptive optics depends on whether atmospheric fluctuations or centroid measurement errors dominate the signal. At the GC the separation between the guide star needed for the adaptive optics infrared wavefront sensor and the star under study is typically $\sim 5''$, corresponding to a separation of 0.25 m at the top (~ 10 km) of the atmosphere. As long as the telescope aperture is larger than this separation, as is the case for Keck and an ELT, the atmosphere dominates and the astrometric precision scales with the telescope diameter as $D^{2/3}$ (Shao & Colavita 1992). A 30 meter ELT is expected to have an astrometric limit that is $3^{2/3} \approx 2$ times smaller than Keck's for $K \lesssim 24$. We therefore adopt $\delta\theta_{30} \sim 0.5$ mas in our calculations, though we consider this a conservative estimate; a 30 meter ELT may attain an astrometric limit as small as 0.1

mas.³

With an adaptive-optics-fed spectrometer on Keck, Ghez et al. (2003a) detected spectral absorption lines in the star S0-2 at a spectral resolution of $R = \lambda/\Delta\lambda \sim 4000$, yielding a radial velocity measurement with an error of 40 km s^{-1} (see also Eisenhauer et al. 2003). Integral field spectroscopy in the near-IR with a 30 meter ELT is expected to enable measurements with $R \sim 1$ – 2×10^4 , suggesting that velocity errors of $\delta v_{30} \sim 10 \text{ km s}^{-1}$ are attainable. This too is a conservative estimate as a 30 meter ELT may achieve velocity errors more than an order of magnitude smaller (D. Figer, private communication). As we discuss below, an ELT will be able to detect stars that are fainter and hence cooler than those currently detectable. Cool stars exhibit rich spectral features including possible molecular lines, enabling high spectral resolution studies. For example, Figer et al. (2003) obtained radial velocities for 85 cool stars in the central parsec of the Galaxy with velocity errors of $\sim 1 \text{ km s}^{-1}$.

Although an ELT's astrometric and spectroscopic limits may differ from the above estimates, we show in § 4.2 that the uncertainties in the model parameters extracted from the monitoring data, such as the distance to the GC and the extended matter profile, scale almost linearly with the measurement errors. The constraints on the parameters for different values of $\delta\theta$ and δv can therefore be readily inferred from our results.

2.2. Confusion Limit

The brighter stars wash out the signal of fainter stars, thereby limiting the luminosity of the faintest observable star. This limit depends on the telescope optics (e.g., angular resolution) and on the stellar luminosity function (LF). Using measurements of stellar photometry near the GC, we now estimate the minimum luminosity that a star at the GC can have and still be identified and monitored with an ELT. For a given star of luminosity l and for a given K -band stellar LF, we determine the integrated flux from all nearby background stars with luminosity $< l$. At some minimum luminosity, the emission from a single star is comparable to the background emission; this luminosity sets the confusion limit.

Following Takeuchi et al. (2001) and references therein, let $x_S = Sh(\theta, \phi)$ be the response of the telescope to a source of flux density S at an angular position (θ, ϕ) from the line-of-sight axis to the center of the source. $h(\theta, \phi)$ is the point-spread function (PSF) of the telescope, normalized to unity at the center. Since all sources at the GC are essentially at the same distance, we can instead express the response in terms of stellar luminosity l , i.e., let $x = lh(\theta, \phi)$. The variance in the telescope response due to crowding is the confusion noise σ . To detect a source with high statistical significance, its luminosity must be greater than some cutoff l_c , or equivalently, x must be greater than a response cutoff x_c . Defining $q = x_c/\sigma$, a source is above the confusion limit if its signal-to-noise $S = x/\sigma > q$, where we take $q = 5$.

If the number of stars per square arcsec with luminosity in the range $(l, l+dl)$ is $dN = \alpha\Phi(l)dl$, where α is the normalization of the LF and $\Phi(l)$ is its shape, then the mean number of

³ For example, see http://tmt/ucolick.org/reports_and_notes/index/htm, Report No. 34.




FIG. 1.— The K-band magnitude limit and number of stars with detectable orbital motions as a function of the aperture of a diffraction limited ELT. Results are shown for power-law K-band luminosity functions normalized to observations by Schödel et al. (2003) with slopes matching the $\sim 2\sigma$ range found by Genzel et al. (2003a). The uncertainty in the number counts is considerably smaller than the uncertainty in the magnitude limits.

source responses of intensity x is

$$\begin{aligned} R(x)dx &= \int \frac{dN(l)}{dl} dl d\Omega \\ &= \int \alpha \Phi \left(\frac{x}{h(\theta, \phi)} \right) \frac{d\Omega}{h(\theta, \phi)} dx, \end{aligned} \quad (2)$$

where the integral is over the solid angle of the PSF. The confusion noise σ due to all sources fainter than x_c is then

$$\sigma^2 = \int_0^{x_c} x^2 R(x) dx. \quad (3)$$

Since we are interested in calculating the cutoff response of a given detector for a given LF, we need to solve for the confusion noise. Assuming a power law LF of the form $dN/dl = \alpha \Phi(l) = \alpha l^{-\eta}$ we have

$$R(x) = \int \alpha \left[\frac{x}{h(\theta, \phi)} \right]^{-\eta} \frac{d\Omega}{h(\theta, \phi)} = \alpha x^{-\eta} \Omega_{\text{eff}}, \quad (4)$$

where

$$\Omega_{\text{eff}} = \int h(\theta, \phi)^{\eta-1} d\Omega. \quad (5)$$

Therefore

$$\sigma^2 = \alpha \Omega_{\text{eff}} \int_0^{x_c} x^{2-\eta} dx = \left(\frac{q^{3-\eta} \alpha \Omega_{\text{eff}}}{3-\eta} \right)^{2/(\eta-1)}. \quad (6)$$

In this paper we only consider power-law LFs, though one can obtain an expression for σ for general forms of the LF (Franceschini et al. 1989).

For a Gaussian PSF $h(\theta, \phi) = h(\theta) = \exp[-(4 \ln 2)(\theta/\theta_0)^2]$, where θ_0 is the PSF's full-width at half-maximum. This gives $\Omega_{\text{eff}} = \pi \theta_0^2 / [(4 \ln 2)(\eta - 1)]$ and thus

$$\sigma = \left(\frac{q^{3-\eta}}{3-\eta} \right)^{1/(\eta-1)} \left[\frac{\pi \theta_0^2 \alpha}{(4 \ln 2)(\eta - 1)} \right]^{1/(\eta-1)}. \quad (7)$$

We now estimate the value of σ for the Keck and an ELT. In the K-band, $\theta_0 \simeq 50$ mas for Keck and $\theta_0 \simeq 15$ mas for a $D = 30$ m ELT. We also need the K-band luminosity function (KLF) of stars at the GC. Genzel et al. (2003a) find that the KLF within $1.5''$ of the GC is well described by a power-law with slope $\beta = d \log N / dK = 0.21 \pm 0.03$ where K is the apparent magnitude in the K-band. We consider KLFs with slopes within the $\sim 2\sigma$ range $0.15 < \beta < 0.27$, which in terms of $\eta = 1 + \beta/0.4$ corresponds to the range $1.38 < \eta < 1.68$.

Schödel et al. (2003) measured the photometry of more than 40 stars in the central arcsec, 29 of which reside within $0.8''$ (~ 6000 AU). We normalize the KLF to these 29 stars. We limit our analysis to these innermost stars since the KLF inside $0.8''$ appears to differ from that outside this region (see § 1). We do not attempt to account for a possible radial dependence but instead assume the KLF is constant.

Of the 29 stars, the brightest has apparent magnitude $K = 13.4$ and the dimmest $K = 17.3$. Assuming a K-band extinction of 3.3 mag (Rieke, Rieke, & Paul 1989) and a distance to the GC $R_0 = 8$ kpc, these apparent magnitudes correspond to K-band luminosities of $l_{\text{min}} = 0.8 L_{\odot}$ and $l_{\text{max}} = 28 L_{\odot}$. For a given η we calculate $\alpha = N_{\text{obs}}(1-\eta)/(l_{\text{max}}^{1-\eta} - l_{\text{min}}^{1-\eta})$ where $N_{\text{obs}} = 29/\pi(0.8'')^2$ and by equation (7) solve for σ . Integrating the luminosity function over stars brighter than $x_c = q\sigma$ yields the number count of detectable stars $N(l > l_c) \sim \alpha x_c^{1-\eta}/(\eta-1)$.

In Figure 1 we show how the K-band magnitude limit and number N of stars with detectable orbital motions (those within 3000 AU of the GC; see below) scale with the aperture of a diffraction limited ELT assuming $\beta = 0.21 \pm 0.06$ ($\eta = 1.53 \pm 0.15$). Since by equation (7) $N \propto x_c^{1-\eta} \propto \theta_0^{-2}$, we find that $N \simeq 100(D/30 \text{ m})^2$. Furthermore, because $x_c \propto \alpha^{1/(\eta-1)}$, $N(l > l_c)$ is not very sensitive to the value of x_c for a fixed η . Therefore, when the above analysis is performed on a subset of the 29 stars within $0.8''$ (e.g., stars within $0.4''$ or alternatively stars brighter than $K = 16$), the derived number counts,

unlike the magnitude limits, do not change significantly. The number counts we derive for an ELT are therefore robust even though the magnitude limits are subject to some uncertainty.

To extract orbital parameters the acceleration of a star in the plane of the sky must be detected, i.e., it must be greater than the threshold acceleration ξ_t . For Keck $\xi_t \sim 1-2 \text{ mas yr}^{-2}$ while for a 30 meter ELT $\xi_t \sim 0.5 \text{ mas yr}^{-2}$. The accelerated proper motion is detectable over the entire orbit if the acceleration at apocenter exceeds the threshold. For a face on orbit this requires $a(1+e) < (GM/\xi_t R_0)^{1/2}$. Thus the acceleration will be detectable with a 30 meter ELT over the entire orbit if $a \lesssim 3000 \text{ AU}$ (period $\lesssim 80 \text{ yr}$). To construct our mock stellar orbits to simulate observations that can be made with an ELT, we only consider orbits satisfying this constraint. As Figure 1 shows, within 3000 AU, approximately 100 stars are brighter than a 30 meter ELT's confusion limit. Furthermore, since the surface density of stars is $\sim 200 \text{ arcsec}^{-2}$, one does not expect to observe stars with apocenters smaller than $\sim 300 \text{ AU}$ ($\sim 0.04''$) with such an ELT. We therefore conclude that a 30 meter ELT will detect the accelerated motion of ~ 100 stars with semi-major axes $\lesssim 3000 \text{ AU}$ and apocenter distances $\gtrsim 300 \text{ AU}$.

Another related issue is the frequency with which an ELT will measure positions and radial velocities for the N monitored stars, given a reasonable commissioning of $\sim 10 \text{ GC}$ exposures per year. As it will be equipped with an integral field unit spectrometer an ELT can obtain simultaneous spectral and spatial data over a relatively large region of sky. It is possible for it to measure positions and velocities for all N stars in a single image. This suggests that a dedicated observing program can reasonably obtain ten measurements per star per year.

At the high levels of precision obtainable with an ELT, orbital parameter constraints should scale with the measurement errors σ (i.e., $\delta\theta$ and δv) and the number of stars N as $\sigma/N^{1/2}$. Based on the above discussion we therefore expect the parameter constraints to scale with telescope aperture as $\sim D^{-2/3}/D \sim D^{-5/3}$. We verify this relation in our numerical simulation results described in § 5.

2.3. Central Point Source — Sgr A*

At radio wavelengths Sgr A* is detected as a non-thermal (Beckert et al. 1996; Serabyn et al. 1997), compact (Rogers et al. 1994), static (Backer & Sramek 1999; Reid et al. 1999), variable (Zhao, Bower, & Goss 2001) source. An X-ray source coincident with Sgr A* has also been detected (Baganoff et al. 2003) and consists of a resolved, steady-state component with size $\sim 1''$ and an unresolved flaring component that increases in flux density by an order of magnitude over the course of a few hours roughly once per day (Baganoff et al. 2001; Goldwurm et al. 2003; Porquet et al. 2003).

Recently, a near-infrared counterpart to Sgr A*, located within a few mas of the dynamically determined black hole position, has been observed in the H band ($1.7\mu\text{m}$; Genzel et al. 2003b) and the L' band ($3.8\mu\text{m}$; Ghez et al. 2004). Like the X-ray emission, the infrared emission consists of a quiescent component and a variable component. The latter exhibits flux densities that increase by a factor of a few over the course of tens of minutes to one week, with possible signs of periodicity (Genzel et al. 2003b). The observed L' -magnitudes are in the range 12.2–13.8, corresponding to L' -luminosities $\sim 10-100L_\odot$.

Although the current sample of stars with detected accel-

erated motion are brighter than the Sgr A* infrared emission, the stars detectable with an ELT will have comparable luminosities. A star that passes near the black hole can therefore be confused with the emission of Sgr A*. Conservatively, such confusion limits monitoring when the projected separation between a star and Sgr A* is smaller than the resolution of the detector. For a 30 meter ELT operating at the diffraction limit in the K band, this corresponds to $\sim 15 \text{ mas}$ (120 AU). However, as we found in § 2.2, confusion with nearby stars precludes such a telescope from detecting orbits with apocenters smaller than 300 AU ($\sim 40 \text{ mas}$) and most of the monitored stars do not therefore pass within 15 mas of Sgr A*. Of those that do, most spend only a small fraction of their total orbital period that close to the black hole; e.g., a star with a semi-major axis of 200 AU and eccentricity 0.9 is within 15 mas of the black hole for only 10% of its orbital period. The same arguments hold for other ELT apertures. Therefore, the infrared emission from the black hole will not significantly impair orbital monitoring with an ELT.

3. ORBITAL DYNAMICS

While current observations of stellar proper motions near the black hole at the GC are consistent with motion around a Newtonian point mass, we show that with an ELT non-Keplerian motions are going to be detectable. There are various effects that cause deviations from Keplerian motion, including the Newtonian retrograde precession (NRP) of an orbit due to the presence of an extended matter distribution (§ 3.2), the relativistic prograde precession (RPP; § 3.3), and the frame dragging effects related to the black hole spin (§ 3.4). In addition, we account for an apparent deformation of the observed proper motion (“Roemer effect”; § 3.5) due to the differences in light travel times at different locations along the orbit. A discussion of the effects of encounters between monitored stars and background stars is given in § 3.6. We now describe the orbital equations of motion and estimate the magnitude of the various non-Keplerian effects. A number of relativistic effects, including those we consider below, are discussed in Pfahl & Loeb (2003) in connection with long-term timing observations of a radio pulsar that might be detected in a $\lesssim 100$ year orbit about the GC.

3.1. Equations of Motion

We found in § 2 that we do not expect a 30 meter ELT to detect orbits with apocenter smaller than $\sim 300 \text{ AU}$ due to confusion noise. Assuming orbits uniformly distributed in e^2 , the probability that a given star has $e > 0.99$ is 2%. Since most of the ~ 100 stars such an ELT monitors will have semi-major axes $> 1000 \text{ AU}$, it is unlikely that any will have pericenter distance smaller than a few AU. As a result, the ratio of the Schwarzschild radius to the pericenter distance of the stars will satisfy $R_s/r_p \lesssim 0.05$, or expressed in terms of the stellar velocity at pericenter, $v_p/c \lesssim 0.2$. The post-Newtonian approximation to the geodesic equations that is accurate to order $(v/c)^2$ provides an adequate description of the stellar orbits given the observational precision expected with an ELT.

The geodesic equation for test particles orbiting a spherically symmetric mass is, in the post-Newtonian approximation (Weinberg 1972; Rubilar & Eckart 2001),

$$\frac{d\mathbf{v}}{dt} = -\nabla\Phi - [2\nabla\Phi^2 + v^2\nabla\Phi - 4\mathbf{v}(\mathbf{v} \cdot \nabla)\Phi - \mathbf{v} \times (\nabla \times \zeta)]/c^2, \quad (8)$$

where $\mathbf{v} = d\mathbf{x}/dt$ is the velocity vector, Φ is a time-independent gravitational potential, and $\zeta = 2G(\mathbf{x} \times \mathbf{J})/r^3$ is a vector potential associated with the spin $\mathbf{J}$ of the gravitating mass, which we assume is constant with time. We assume the density distribution of the extended matter at radii traversed by the stars and smaller is a power-law profile $\rho(r) = \rho_0(r/r_0)^{-\gamma}$. Input model parameters are described in § 4.2.

The relativistic effects include corrections to the orbital dynamics and to the observed motion due to propagation effects. The former class includes the RPP and frame dragging while the later class includes the lowest order (v/c) Roemer time delay and such $(v/c)^2$ effects as time dilation, gravitational redshift, and the Shapiro time delay. Since the $(v/c)^2$ propagation effects each have different functional dependences on the orbital parameters (see e.g., Pfahl & Loeb 2003), including them may break degeneracies, though they may also weaken the sensitivity to some parameters. However, our interest in the $(v/c)^2$ relativistic effects is primarily connected with the ability of an ELT to probe general relativity on the scale of a massive black hole rather than with the effects' potential utility for parameter estimation. Since $v_p/c < 0.2$, relativistic effects help to constrain the orbital parameters by at most a few percent. We therefore chose not to include the $(v/c)^2$ propagation effects in our analyses. The analysis of actual data obtained from an ELT must, however, account for all the relativistic effects.

3.2. Newtonian Retrograde Precession

The NRP was discussed in the context of the GC by Rubilar & Eckart (2001). An extended matter distribution causes stellar orbits to precess due to differences in the amount of mass that is contained between the apocenter and the pericenter radii. In the numerical calculations that follow, we determine how much an orbit precesses due to the extended matter by solving equation (8). Here, however, we obtain an estimate of the magnitude of the precession for stars at the GC by considering the potential of the extended matter to be a small correction $\delta\Phi$ to the potential of the black hole. Expanding the total potential to linear order in $\delta\Phi$, the angular shift per period is (Landau & Lifshitz 1960)

$$\Delta\phi_{\text{Newt}} = \frac{\partial}{\partial L} \left(\frac{2}{L} \int_0^\pi r^2 \delta\Phi d\varphi \right), \quad (9)$$

where L is the orbital angular momentum per unit mass, $r = a(1 - e^2)/(1 + e \cos \varphi)$, and φ is the phase of the orbit. If $\gamma < 2$ we have $\delta\Phi = \beta r^{2-\gamma}$, where $\beta = GM_{\text{ext}}/(2 - \gamma)r_0^{3-\gamma}$ is a constant and $M_{\text{ext}} \equiv M_{\text{ext}}(r < r_0)$ is the extended matter mass within r_0 . The orbital precession is then given by (Munyanza & Viollier 1999)

$$\Delta\phi_{\text{Newt}} = \frac{2\beta}{GM_{\text{bh}}} [a(1 - e^2)]^{3-\gamma} g(\gamma, e), \quad (10)$$

where

$$g(\gamma, e) = \frac{1 - e^2}{e^2} (4 - \gamma) [I_{4-\gamma}(e) - I_{5-\gamma}(e)] + (7 - 2\gamma) I_{4-\gamma}(e) \quad (11)$$

and

$$I_n(e) \equiv \int_0^\pi \frac{d\varphi}{(1 + e \cos \varphi)^n}. \quad (12)$$

Assume that the extended matter consists of stars with $\gamma = 7/4$ and $\rho_{1\text{pc}} = 2 \times 10^5 M_\odot \text{ pc}^{-3}$ (see § 1). Consider an

S0-2 like orbit with a semi-major axis of 0.005 pc and eccentricity $e = 0.9$. The enclosed stellar mass at apocenter and pericenter are $6000 M_\odot$ and $150 M_\odot$. Solving equation (10) yields a precession per revolution of $\Delta\phi_{\text{Newt}} \approx 0.08^\circ$, corresponding to an apparent angular apocenter shift of roughly $\Delta\phi_{\text{Newt}} a(1 + e)/R_0 \approx 0.3 \text{ mas}$. Thus, a few S0-2 like orbits with astrometric errors of 0.5 mas provide a meaningful constraint on the stellar distribution within the inner few milliparsecs. If the density of the dark matter cusp at the stellar positions exceeds $\sim 10^8 M_\odot \text{ pc}^{-3}$, then it too will produce a detectable precession; it will not be easily distinguished from the stellar contribution (however see § 3.6).

3.3. Relativistic Prograde Precession

The RPP causes a pericenter advance per revolution of (see Weinberg 1972) $\Delta\phi_{\text{pro}} = 3\pi R_s/a(1 - e^2)$, where $R_s = 2GM_{\text{bh}}/c^2$ is the Schwarzschild radius of the black hole. The magnitude of the effect is $\propto (v/c)^2$. The apparent apocenter shift per revolution caused by the RPP is $\Delta s \approx \Delta\phi_{\text{pro}} a(1 + e)/R_0 = 3\pi R_s/R_0(1 - e)$, which corresponds to an apparent shift of $\sim 1 \text{ mas}$ for the star S0-2. Although the RPP has an additional factor of (v/c) relative to the Roemer effect (§ 3.5), this attenuation can be compensated by having a few high eccentricity stars in the sample. Furthermore, unlike the Roemer effect, the RPP shift is to first order independent of the semi-major axis and is therefore equally sensitive to stars at all radii (although stars at large radii also have long periods). Consider an orbit seen face on and observed for N_{orb} complete periods. Since the precession angles per revolution add linearly, the signal-to-noise from the RPP is $S_{\text{pro}} \sim \Delta s N_{\text{orb}}/\delta\theta$, or

$$S_{\text{pro}} \sim 0.1 \frac{N_{\text{orb}}}{1 - e} \left(\frac{M_{\text{bh}}}{4 \times 10^6 M_\odot} \right) \left(\frac{R_0}{8 \text{ kpc}} \right)^{-1} \left(\frac{\delta\theta}{1 \text{ mas}} \right)^{-1}. \quad (13)$$

In a sample of 100 stars observed with astrometric errors of 0.5 mas and having an eccentricity distribution uniform over e^2 , we expect on average eight stars with $e > 0.96$. If only one such a star is followed over just a single period, the RPP shift will be measured to 5- σ accuracy.

3.4. Frame Dragging

For a spinning black hole, frame dragging effects also cause a precession of the pericenter. The spin precession per revolution for a star orbiting a black hole with spin angular momentum J is given approximately by (see Weinberg 1972, equation (9.5.22); note different notation)

$$\begin{aligned} \Delta\phi_{\text{spin}} &\approx -8\pi j \left(\frac{GM_{\text{bh}}}{cL} \right)^3 \cos\psi \\ &= -\frac{2\sqrt{2}j\Delta\phi_{\text{pro}}}{3} \sqrt{\frac{R_s}{a(1 - e^2)}} \cos\psi, \end{aligned} \quad (14)$$

where ψ is the angle between the orbital angular momentum vector and the black hole spin axis and $0 \leq j \equiv cJ/GM_{\text{bh}}^2 \leq 1$ is the black hole spin parameter.

The black hole spin induces an apocenter shift that is smaller than the RPP shift by a factor of $\sim v/c$. Even if the black hole is maximally spinning ($j = 1$), the shift represents only a 5% contribution on top of the RPP for a star with $a = 200 \text{ AU}$ and eccentricity $e = 0.92$. For an orbit observed face-on the signal-to-noise from a spin-induced apoc-

enter shift is

$$S_{\text{spin}} \approx \frac{2\pi\sqrt{2}j}{R_0\sqrt{a(1+e)}} \left(\frac{R_s}{1-e}\right)^{3/2} \frac{N_{\text{orb}}}{\delta\theta} \cos\psi$$

$$\approx 0.001 \frac{jN_{\text{orb}} \cos\psi}{\sqrt{(1+e)(1-e)^3}} \left(\frac{M_{\text{bh}}}{4 \times 10^6 M_\odot}\right)^{3/2}$$

$$\times \left(\frac{a}{1000 \text{ AU}}\right)^{-1/2} \left(\frac{R_0}{8 \text{ kpc}}\right)^{-1} \left(\frac{\delta\theta}{1 \text{ mas}}\right)^{-1}. \quad (15)$$

For example, a 5- σ detection is achieved with an ELT with $\delta\theta = 0.5$ mas if a star with $a = 300$ AU and $e = 0.99$ is monitored for three complete orbits. We expect a 30 meter ELT to detect one star with a semi-major axis that small (§ 2). Assuming eccentricities uniformly distributed in e^2 the probability that star has $e > 0.99$ is only $\sim 2\%$. If $\delta\theta = 0.05$ mas, a star with $a = 300$ AU and $e = 0.95$ will yield a 5- σ detection after being monitored for three complete orbits. Since so high a resolution requires an ELT with aperture $D \sim 100$ m, there will be several stars with $a \lesssim 300$ AU and of these $\sim 10\%$ will have $e > 0.95$. Detecting such a stellar orbit is therefore not unlikely. The spin-induced orbital precession thus requires an astrometric precession of $\delta\theta \lesssim 0.05$ mas (see also Jaroszynski 1998; Fragile & Mathews 2000).

3.5. The Roemer Time Delay

For orbits with non-zero inclination, the distance between the Earth and star, and hence the difference in time between stellar emission and observation, varies with orbital phase. This time delay, given by $\Delta t = t_{\text{obs}} - t_{\text{em}} = z(t_{\text{em}})/c$, where $z(t)$ is the relative distance between the star and the massive black hole, was first recognized by Roemer in 1676 in application to the phases of Jupiter's moons. Unlike the relativistic Doppler effect which includes corrections of order $(v/c)^2$ and higher, the Roemer delay is the classical Doppler effect which only includes terms up to order v/c (see, e.g., Loeb 2003). The delay has a magnitude corresponding to a few percent of a year for an S0-2 like orbit, and is observed as an additional shift in the apparent stellar position with time, $\Delta s(t)$. For a circular orbit seen edge-on the stellar positions $z(t)$ and $s(t)$ are sinusoidal so that

$$\Delta s(t)/a = \cos(\omega t_{\text{obs}}) - \cos(\omega t_{\text{em}})$$

$$= \cos[\omega(t_{\text{em}} + a/c \sin(\omega t_{\text{em}}))] - \cos(\omega t_{\text{em}})$$

$$\simeq -\frac{\omega a}{c} \sin^2(\omega t_{\text{em}}), \quad (16)$$

where $\omega = 2\pi/P$ and we used the fact that for orbits at the GC $v \ll c$. The maximum shift, in units of the semi-major axis, is therefore v/c . For non-zero eccentricity and arbitrary inclination the star's projected position and distance as a function of time are (see e.g., Murray & Dermott 1999)

$$\begin{Bmatrix} x(t) \\ y(t) \\ z(t) \end{Bmatrix} = r(t) \begin{Bmatrix} \cos[\varphi(t) + \alpha] \\ \sin[\varphi(t) + \alpha] \cos i \\ \sin[\varphi(t) + \alpha] \sin i \end{Bmatrix} \quad (17)$$

where i is the inclination, $\varphi(t)$ is the orbital-phase (i.e., the true anomaly), α is the argument of pericenter, and we chose the reference direction so that the x -axis coincides with the longitude of ascending node. The Roemer shift is then $|\Delta s| = (\Delta x^2 + \Delta y^2)^{1/2}$ where $\Delta x = x(t_{\text{obs}}) - x(t_{\text{em}})$ and similarly for Δy . To linear order in v/c the orbit-averaged Roemer shift can be written as

$$\langle |\Delta s(t)|/a \rangle \simeq \frac{\omega a}{c} \sin i \sqrt{f(e, i, \alpha)} \quad (18)$$

where $f(e, i, \alpha)$ is a factor of order unity. For the two extreme cases $\alpha = 0$ and $\alpha = \pi/2$ (corresponding to the line-of-node along the major-axis and minor-axis, respectively), $f(e, i, \alpha)$ is given by

$$f(\alpha = 0) = \frac{\sqrt{1-e^2}}{2e^4} \left\{ 2e^4 \cos^2 i \left(3 - \sqrt{1-e^2} \right) \right.$$

$$+ 3e^2 \left[2 - \sqrt{1-e^2} - \cos^2 i \left(4 - 3\sqrt{1-e^2} \right) \right]$$

$$\left. - 6\sin^2 i \left(1 - \sqrt{1-e^2} \right) \right\}, \quad (19)$$

$$f(\alpha = \pi/2) = \frac{\sqrt{1-e^2}}{2e^4} \left\{ -2e^4 \left(2 - \sqrt{1-e^2} \right) \right.$$

$$+ e^2 \left[10 - 7\sqrt{1-e^2} - \cos^2 i \left(4 - \sqrt{1-e^2} \right) \right]$$

$$\left. - 6\sin^2 i \left(1 - \sqrt{1-e^2} \right) \right\}. \quad (20)$$

An ELT will be able to detect the effect of the Roemer delay in orbits at the GC. The signal-to-noise from N_{obs} observations of an orbit measured with astrometric errors $\delta\theta$ is approximately $S_{\text{delay}} \sim \langle \Delta s \rangle N_{\text{obs}}^{1/2} / R_0 \delta\theta$, or

$$S_{\text{delay}} \approx 0.8 \sqrt{N_{\text{obs}}} \sin i \sqrt{f(e, i, \alpha)} \left(\frac{a}{1000 \text{ AU}} \right)^{1/2}$$

$$\times \left(\frac{M_{\text{bh}}}{4 \times 10^6 M_\odot} \right)^{1/2} \left(\frac{R_0}{8 \text{ kpc}} \right)^{-1} \left(\frac{\delta\theta}{1 \text{ mas}} \right)^{-1} \quad (21)$$

If, e.g., we pick $i \sim \pi/3$, $\alpha = 0$, and $e \sim 1/\sqrt{2}$, an astrometric error of 0.5 mas, a mean semi-major axis of 1000 AU, and 10 observations per star, then we can detect the delay to $S_{\text{delay}} \sim 5$ with roughly 10 stars. We therefore expect the Roemer delay to be detectable in such an ELT's sample of ~ 100 stars and the effect must be taken into account during parameter estimation.

3.6. Interstellar Interactions

In the previous sections we described the motion of a star in the potential of a black hole and a smooth distribution of extended matter, including stars, remnants, and dark matter. This approximation ignores the fact that the potential due to stars and remnants is the sum of discrete point-mass potentials and is therefore not perfectly smooth. The stars experience perturbations due to nearby encounters with individual stars and due to fluctuations in the potential arising from all stars. These perturbations cause a star's orbital parameters to change with time. The magnitude and the rate of these changes depend on the stellar mass function since the perturbations are sensitive to the characteristic mass of the field stars. Thus, measuring the effects of stellar encounters is a probe of the mass function in the central parsec. It also breaks the degeneracy between the contributions of stellar matter and dark matter to the Newtonian orbital precession. Encounters may also be a source of noise in measurements of orbital parameters such as the black hole mass and distance to the GC. While we do not include the effects of encounters in our numerical calculations presented in § 5, we now estimate their magnitude and demonstrate that the encounters might be detectable with an ELT and present a powerful probe of the mass function of stellar remnants at the GC.

An encounter between a test star of mass m_j and a field star of mass m_i with impact parameter b induces a change in the

test star velocity given by (see, e.g., Spitzer 1987)

$$\delta v = \frac{2m_i v_{\text{rel}}}{m_i + m_j} \left[1 + \left(\frac{b}{b_0} \right)^2 \right]^{-1/2}, \quad (22)$$

where $b_0 = G(m_i + m_j)/v_{\text{rel}}^2$ and v_{rel} is the initial relative velocity of the stars. The encounter induces a change in the test star's velocity distinct from that due to orbital motion around the black hole. We solve for the maximum impact parameter b_{max} such that an encounter induces a change in velocity of the test star larger than the minimum detectable change δv_{min} . For uncorrelated position measurements the minimum detectable change in velocity is $\delta v_{\text{min}} \sim \delta \theta R_0 / \sqrt{N_{\text{obs}}} T$, where T is the time baseline over which the orbit is monitored, and N_{obs} is the number of position measurements taken in time T . Assuming $\delta \theta = 0.5$ mas (§ 2.1), $T = 10$ yr, and $N_{\text{obs}} = 100$ yields $\delta v_{\text{min}} \sim 0.2 \text{ km s}^{-1}$. By equation (22) we have

$$b_{\text{max}} = b_0 \sqrt{\left(\frac{2m_i}{m_i + m_j} \frac{v_{\text{rel}}}{\delta v_{\text{min}}} \right)^2 - 1} \approx \frac{2Gm_i}{v_{\text{rel}} \delta v_{\text{min}}}, \quad (23)$$

where the approximation assumes $v_{\text{rel}} \gg \delta v_{\text{min}}$ and $m_j \lesssim m_i$. Assume $v_{\text{rel}} \sim v_p = (GM(1+e)/(1-e)a)^{1/2}$ for an encounter near pericenter and $v_{\text{rel}} \sim (GM/a)^{1/2}$ for an encounter near apocenter. If we take $\delta v_{\text{min}} \sim 0.2 \text{ km s}^{-1}$ and $m_i = 10M_{\odot}$ then for an S0-2 like orbit, $b_{\text{max}} \sim 10 \text{ AU}$ at pericenter and $b_{\text{max}} \sim 50 \text{ AU}$ at apocenter.

We ignore the effect of the black hole on the encounter and treat the interaction between the stars as a two-body problem. This is a fair approximation as long as the duration of the encounter is much shorter than the time scale over which the orbital velocity changes significantly due to the influence of the black hole. At pericenter passage, where the orbital acceleration is greatest, the orbital time scale is $t_p \sim (1-e)^{3/2}P$, where P is the orbital period. The two-body approximation is valid as long as the duration of the encounter satisfies $t_{\text{enc}} \sim b_{\text{max}}/v_{\text{rel}} \ll t_p$. For an S0-2 like orbit $t_p \sim 0.5 \text{ yr}$ while by equation (23) $t_{\text{enc}} \lesssim 0.01 \text{ yr}$ even for $m_i = 20M_{\odot}$.

Next, we estimate the rate at which encounters $b < b_{\text{max}}$ occur for a star on a given orbit (see, e.g., Yu 2003). Let $\Gamma_{ij}(\mathbf{r}, \mathbf{v}_j, t) dm_i$ be the rate at which a star with mass m_j at position $\mathbf{r}$ with velocity $\mathbf{v}_j$ at time t encounters stars with masses in the range $m_i \rightarrow m_i + dm_i$. Assume the number density of stars is spherically symmetric and follows a power law $\nu(r) = \nu_0(r/r_h)^{-\alpha}$. The phase-space distribution function of the stars is given by (Magorrian & Tremaine 1999) $f(\mathcal{E}) = h(\alpha)\mathcal{E}^{\alpha-3/2}$, where $\mathcal{E} = \Psi(r) - v^2/2$ and $\Psi(r)$ is the relative gravitational potential at r , while

$$h(\alpha) = (2\pi\sigma_h^2)^{-3/2}\nu_0 \frac{\Gamma(\alpha+1)}{\Gamma(\alpha-1/2)} \sigma_h^{-2\alpha+3}, \quad (24)$$

with σ_h the linear stellar velocity dispersion outside the sphere of influence of the BH $r_h \simeq 1 \text{ pc}$. The rate of detectable encounters in the mass bin is then

$$\begin{aligned} \Gamma_{ij} &= \int_0^\infty dv_i \frac{2\pi v_i}{v_j} f(\mathbf{r}, v_i) \int_{|v_i - v_j|}^{v_i + v_j} dv_{\text{rel}} v_{\text{rel}}^2 \Sigma(v_{\text{rel}}) \\ &= 2\pi K h(\alpha) \\ &\quad \times \int_0^{\sqrt{2\Psi}} \left(\Psi - \frac{v_i^2}{2} \right)^{\alpha-3/2} \frac{v_i}{v_j} (v_i + v_j - |v_i - v_j|) dv_i, \end{aligned} \quad (25)$$

where the cross section for detectable encounters $\Sigma = \pi b_{\text{max}}^2$, and $K = 4\pi G^2 m_i^2 / \delta v_{\text{min}}^2$.

We now determine the rate at which stars that will be monitored with an ELT undergo detectable encounters. The integral in equation (25) is most easily evaluated in the special case $\alpha = 3/2$, which is compatible with current observational constraints (Genzel et al. 2003a). To obtain a rough estimate of the rates, consider the case $\alpha = 3/2$ and assume the background stars all have identical mass not smaller than that of the test star (e.g., they are a population of stellar mass black holes). By equation (25)

$$\Gamma_j(\mathbf{r}, \mathbf{v}_j, t) = \frac{3\sqrt{2}}{8} \frac{K\nu_0}{\sigma_h^3} (2\Psi - \frac{1}{3}v_j^2), \quad (26)$$

and upon averaging over the orbital phase

$$\begin{aligned} \Gamma_j(e, a) &= \frac{1}{P} \int_0^P \Gamma_j(\mathbf{r}, \mathbf{v}_j, t) dt \\ &= \frac{5\pi\sqrt{2}\nu_0}{2\sigma_h^3} \left(\frac{Gm_i}{\delta v_{\text{min}}} \right)^2 \frac{GM_{\text{bh}}}{a}. \end{aligned} \quad (27)$$

Assume the N stars monitored with an ELT have an eccentricity distribution uniform in e^2 (isotropic velocity ellipsoid) so that $dN/de da \propto ea^{2-\alpha} = e\sqrt{a}$. Integrating over these distributions and normalizing to $N = N(< a_2) \propto a_2^{3/2}$ yields the total rate at which encounters are detected with an ELT

$$\begin{aligned} \Gamma(\alpha = 3/2) &= \int_0^1 \int_{a_1}^{a_2} \frac{dN(e, a)}{de da} \Gamma_j(e, a) de da \\ &= \frac{15\pi\sqrt{2}\nu_0}{2\sigma_h^3} \left(\frac{Gm_i}{\delta v_{\text{min}}} \right)^2 \frac{GM_{\text{bh}}N}{a_1 + a_2 + \sqrt{a_1 a_2}}, \end{aligned} \quad (28)$$

where a_1 and a_2 define the range in semi-major axis that is accessible to observations.

Given the above expression for the encounter rate for $\alpha = 3/2$, we rely on scaling relations to estimate the rate for different α . Since the encounter rate is proportional to the stellar density, $\Gamma(\alpha) \simeq \Gamma(3/2)(r_h/r)^{\alpha-3/2}$. Thus, if $\alpha = 7/4$, the rate of encounters is ~ 3 times larger than for $\alpha = 3/2$. The time scale for detectable encounters is therefore

$$\begin{aligned} \Gamma^{-1} &\sim 0.3 \text{ yr} \left(\frac{m_i \nu_0}{2 \times 10^5 M_{\odot} \text{ pc}^{-3}} \right)^{-1} \left(\frac{m_i}{10 M_{\odot}} \right)^{-1} \\ &\quad \times \left(\frac{N}{100} \right)^{-1} \left(\frac{a_2}{3000 \text{ AU}} \right) \left(\frac{\sigma_h}{100 \text{ km s}^{-1}} \right)^3 \\ &\quad \times \left(\frac{\delta \theta}{0.5 \text{ mas}} \right)^2 \left(\frac{T}{10 \text{ yr}} \right)^{-2} \left(\frac{N_{\text{obs}}}{100} \right) \left(\frac{a_2}{r_h} \right)^{\alpha-3/2}, \end{aligned} \quad (29)$$

where we use the results of § 2 that $N \approx 100$, $a_1 \simeq 200 \text{ AU}$, and $a_2 \simeq 3000 \text{ AU}$, and have also assumed that the mass density of background particles $m_i \nu_0$ is constant and independent of m_i .

Therefore, assuming a density cusp dominated by $\sim 10M_{\odot}$ black holes, ~ 30 nearby stellar encounters will be detectable during ten years of monitoring with an ELT with $\delta \theta = 0.5 \text{ mas}$. Measurement of the frequency of detectable orbital deflections $\Gamma \propto m_i$ is a direct test of the average mass of the dark remnants that probably dominate the mass density near the black hole (Morris 1993; Miralda-Escudé & Gould 2000) but are otherwise not directly detectable. Since $N(< a) \propto a^{3/2}$,

then by equation (28), $\Gamma \propto a^{1/2}$, i.e., the encounter rate increases with distance from the massive black hole. The stars at $a > 3000$ AU with detectable linear proper motion may therefore yield the strongest constraint on the mass function of stellar remnants, despite being below the threshold for detecting accelerated motion due to the massive black hole.

4. METHOD

In this section we describe how we generate mock ELT orbital data. We also describe our implementation of the Markov Chain Monte Carlo (MCMC) method which we use to estimate the uncertainties in the orbital parameters and map the shape of the likelihood surface.

4.1. Parameter Estimation

We are interested in estimating the uncertainties in the parameters given proper motion and radial velocity information for a sample of N stars orbiting the massive black hole at the GC. Each star's projected orbit is described by six phase space parameters. The black hole mass, its 3-dimensional position, and the normalization and slope of the extended matter distribution, contribute an additional six parameters. The entire parameter space of our model therefore has dimension $J = 6N + 6$.

Parameter estimation on a J -dimensional grid is not practical. Since the computational cost of the grid-based approach increases exponentially with J , the parameter space becomes prohibitively large for even just two or three stars. By contrast, the cost of the MCMC method scales almost linearly with J .

We now briefly describe the basic ideas of the MCMC method and our choice of implementation. A general discussion of the theory and application of the MCMC approach is given in Gilks, Richardson, & Spiegelhalter (1996). Readers not interested in the details of our parameter estimation scheme can skip ahead to § 4.2.

Let D denote the observed data, θ the model parameters, $P(\theta)$ the prior distribution (which is uniform here), and $L(D|\theta)$ the likelihood of detecting the data for a given set of parameter values. By Bayes's theorem the distribution of θ conditioned on D is given by

$$\pi(\theta|D) = \frac{P(\theta)L(D|\theta)}{\int P(\theta)L(D|\theta)d\theta}, \quad (30)$$

and is called the posterior distribution of θ . The statistical properties of the parameters such as means, moments and confidence contour levels, are entirely specified by $\pi(\theta|D)$.

Explicit evaluation of the integral in the denominator of equation (30) is not practical in large dimensional models. The MCMC method avoids evaluating the integral by instead generating a Markov chain of parameter points the distribution of which converges to the posterior distribution $\pi(\theta|D)$. The Markov aspect refers to the property that the probability distribution of the n th state (i.e., point) in the chain θ_n depends only on the previous state θ_{n-1} . It can be shown (e.g., Gilks, Richardson, & Spiegelhalter 1996) that the density of points in a Markov chain converges to $\pi(\theta|D)$ if the following criteria are satisfied: (1) the chain is irreducible, namely from any starting state θ_0 the chain can reach any non-empty set with positive probability in some finite number of iterations; (2) the chain is aperiodic in that it does not oscillate between different sets of states in a regular periodic fashion; (3) the chain is positive recurrent, meaning that if the initial

value θ_0 is sampled from the posterior then the expected time (i.e., number of iterations) to return arbitrarily close to state θ_0 is finite. There are several algorithms for generating Markov chains that satisfy the above properties. We use the Metropolis algorithm (Metropolis et al. 1953) in our numerical calculations.

Our implementation of the Metropolis algorithm is as follows.

1. Start a chain at $t = 0$ with some initial state θ_0 .
2. Generate a trial state θ' according to the jump proposal distribution $q(\theta'|\theta_t)$ (see below). Compute

$$\alpha(\theta_t, \theta') = \min \left[1, \frac{L(D|\theta')}{L(D|\theta_t)} \right]. \quad (31)$$
3. Sample a uniform random variable U that lies between $(0, 1)$.
4. If $U \leq \alpha(\theta_t, \theta')$ then set $\theta_{t+1} = \theta'$ (i.e., accept the jump). If $U > \alpha(\theta_t, \theta')$ then set $\theta_{t+1} = \theta_t$.
5. Increment t .
6. Go to step #2.

If the observational errors follow a normal distribution, $L(D|\theta) \simeq \exp[-\chi^2(\theta, D)/2]$. The $\chi^2(\theta, D)$ statistic for a single star is given by

$$\chi^2(\theta, D) = \sum_{i=1}^M \left\{ \frac{[x_i(\theta) - x_i(D)]^2}{\sigma_{x,i}^2} + \frac{[y_i(\theta) - y_i(D)]^2}{\sigma_{y,i}^2} \right\} + \sum_{j=1}^K \frac{[v_j(\theta) - v_j(D)]^2}{\sigma_{v,j}^2}, \quad (32)$$

where (x, y) is the astrometric position of the star, v its radial velocity, and σ the corresponding measurement errors (i.e., $\sigma_{x,y} = \delta\theta$ and $\sigma_v = \delta v$). We simultaneously fit to multiple stars by summing each star's χ^2 to form a cumulative χ^2 for the model.

The jump proposal distribution $q(\theta'|\theta_t)$ is the probability of selecting a trial state θ' given the current state θ_t . For the Metropolis algorithm one considers only symmetric proposals of the form $q(\theta'|\theta_t) = q(\theta_t|\theta')$. We choose to model the jump distribution as a multivariate normal distribution with mean θ_t and constant covariance matrix C .

Although the distribution of points in a chain is independent of the form of the jump distribution once the Markov chain has converged, the time it takes a chain to converge is sensitive to the jump distribution. To ensure an efficient run one must carefully choose the shape and step size of the jump distribution. An ideal jump distribution has a shape and step size that not only minimizes the convergence time but also samples the entire posterior distribution efficiently. In our implementation the shape of the jump distribution is determined by C and the step size is determined by a constant scale factor multiplying C .

C is chosen such that the shape of the jump distribution is similar to that of the posterior distribution, although we again emphasize that the shape is only important for the efficiency of convergence. This ensures that the chain mixes well even in regions of degeneracy. To this aim, we compute the covariance matrix that describes the shape of the χ^2 surface in

the neighborhood of its minimum. We first compute the approximate best fit parameter state θ_{bf} by minimizing χ^2 . We then specify a pilot covariance matrix C_p that is purely diagonal with variances given by a reasonable guess of the 1σ uncertainties for individual parameters. We draw a number (~ 1000) of pilot points from a multivariate normal distribution with mean θ_{bf} and covariance C_p . Since the pilot points are within $\sim 1\sigma$ of the χ^2 minimum, the shape of the χ^2 surface in the region of the points is approximately quadratic. We solve the linear least-squares problem by fitting a quadratic χ^2 model to the points and obtain the approximate Fisher matrix that describes the curvature of the χ^2 surface. We then determine the eigenvalues and eigenvectors of the Fisher matrix. If any of the eigenvalues are negative, indicating that the shape of the surface is unconstrained in some direction, we generate more pilot points and redo the linear least-squares fit. Finally, we invert the resulting Fisher matrix to obtain the covariance matrix C .

The constant scale factor that determines the step sizes must also be carefully chosen to ensure that the chain effectively explores the parameter space. If the steps are too small the chain does not mix well as it stays in one region of the parameter space for long periods of time. If the steps are too large, the trial states are rejected frequently. For a multivariate normal jump distribution the most efficient step sizes are those for which $\sim 25\%$ of the jump proposals are accepted (see Gelman 1995). We chose the (constant) jump scale factor to optimize the acceptance rate.

The first steps in a chain may be sensitive to the starting state θ_0 and are therefore not sampled from the posterior distribution. We discard these initial “burn in” points. To ensure that a chain has converged and is sampling the full posterior distribution we run multiple chains each starting at widely dispersed states. We tested for convergence with the Gelman-Rubin test statistic (Gilks, Richardson, & Spiegelhalter 1996).

4.2. Mock Data

To generate a realistic set of orbital data we must determine: (i) the number of stars N we can detect and monitor with an ELT, (ii) the spatial distribution of these stars, (iii) the number of observations per year per star and, (iv) the observational errors in the stellar positions and velocities. In § 2 we showed that with a 30 meter ELT the position of the stars can be centroided to an astrometric precision $\delta\theta_{30}$ between $0.1 - 0.5$ mas and the radial velocities measured to accuracies δv_{30} between $1 - 10$ km s $^{-1}$. We found that with such an ELT we can detect the accelerated proper motion of approximately 100 stars. We estimate that an integral-field spectrograph on an ELT enables a dedicated GC observing program to obtain position and velocities of each of the 100 stars roughly ten times per year.

A realistic mock data set might therefore consist of $N = 100$ stars, observed over a ten year baseline with ten observations per year per star, with position and velocity measurements for each star accurate to 0.5 mas and 10 km s $^{-1}$. Unfortunately, running our MCMC simulation on such a large data set was not feasible due to limits in computational speed. A typical run requires $\sim 10^7$ iterations (i.e., jumps) in order to fully sample the posterior distribution. This corresponds to a minimum of ~ 3 days on a desktop machine for just 20 stars ($J \simeq 126$) with 100 points per star; a simulation with 100 stars takes approximately five times longer. However, one can obtain realistic results from a reduced sample size by properly

TABLE 1
STELLAR ORBITAL PARAMETERS

Star	P (yr)	a (AU)	e	r_{min} (AU)	i (deg)
1	2.9	325	0.852	48	89
2	5.2	476	0.587	197	104
3	5.4	488	0.783	106	46
4	9.5	712	0.560	313	86
5	13.4	895	0.800	179	114
6	14.2	931	0.772	212	81
7	17.7	1078	0.800	216	15
8	23.2	1289	0.605	510	126
9	25.2	1363	0.903	132	25
10	26.6	1413	0.730	382	142
11	36.8	1755	0.705	517	56
12	40.4	1869	0.890	206	93
13	42.3	1928	0.885	223	105
14	46.6	2056	0.703	611	42
15	50.6	2172	0.552	973	104
16	51.5	2197	0.567	951	145
17	52.2	2218	0.439	1245	106
18	52.5	2225	0.636	810	95
19	57.8	2372	0.319	1617	26
20	78.1	2901	0.220	2264	60

NOTE. — The listed parameters are: orbital period (P), semi-major axis (a), eccentricity (e), pericenter distance (r_{min}), and inclination (i). They are not all independent variables.




FIG. 2.— Astrometric positions of the 20 synthesized orbits to which we fit the model. The thick-lined portion of each orbit is the proper motion over the fitted 10 year baseline assuming 10 observations per year.

scaling the χ^2 values (see equation 32) to emulate the full sample size. In particular, we construct a mock data set with $N = 20$ and multiply the χ^2 of each star by a factor of five.

This approach yields realistic estimates of parameter uncertainties as long as the mock data set with $N = 20$ stars fairly represents the full data set with 100 stars. We minimize the effects of sample variance as follows. We first generate data for 1000 synthetic orbits. These orbits are drawn from the

distribution function of the power-law density profile assuming randomly oriented orbits and considering only those orbits with semi-major axes in the range detectable with an ELT (see § 2). We generate mock data for these orbits assuming Gaussian position and velocity errors with dispersions $\delta\theta$ and δv and a specific input model for the potential (e.g., black hole plus extended matter). For each individual star, we compute the difference in χ^2 between the best-fit model (essentially the model used to generate the data) and the null hypothesis model (e.g., no extended matter). We then rank the stars by the size of this χ^2 difference. We bin the 1000 stars into N bins according to their rank and randomly select one star from each bin. The resulting N stars form the set of orbits to which we fit.

Table 1 lists the orbital parameters for one realization of a sample of 20 stars to which we fit. Given the orbital parameters, we generate mock data by solving the equation of motion for each star (see equation 8). In Figure 2 we show the astrometric positions of the 20 stars over the ten year observational baseline with ten epochs per year. The values of the input model parameters describing the potential are: $M_{\text{bh}} = 4 \times 10^6 M_\odot$, $R_0 = 8$ kpc, $(x_{\text{bh}}, y_{\text{bh}}) = (0, 0)$, $M_{\text{ext}}(r < 0.01 \text{ pc}) = 6000 M_\odot$ and either $\gamma = 1.5$ or 2.

To test that the parameter uncertainty estimates are not affected by sample variance we ran simulations on several different draws of 20 stars. As we show in § 5 the parameter uncertainties obtained are similar amongst the different data sets, suggesting that sample variance does not affect the results. Thus, given the current uncertainties in an ELT's ultimate capabilities as well as the uncertainty in the exact nature of the stellar distribution at the GC, we conclude that to a reasonable approximation a mock data set comprised of $N = 20$ stars with χ^2 values increased five-fold yields parameter uncertainties similar to that expected with observations by a 30 meter ELT.

We also show in the next section that the orbital parameter constraints scale with the measurement errors σ and number of stars N as $\sigma/N^{1/2}$. Results for a wide range of assumed ELT capabilities (i.e., different $\delta\theta$ and δv , different aperture, etc.) can therefore be computed by scaling the results of our fiducial 30 meter ELT model, using the relations between N , aperture D , $\delta\theta$, and δv , given in § 2.

5. RESULTS

In this section we investigate how well observations with an ELT constrain the structure of the GC. Our model of the GC and the orbits was described in § 3. We draw stellar orbital parameters from a phase-space distribution determined by the model and use these orbits to synthesize mock ELT data (see § 4). We then fit a model to the mock data and calculate the uncertainties in the parameters using the MCMC technique discussed in § 4.1. We show results for a 30 meter ELT with $(\delta\theta, \delta v) = (0.5 \text{ mas}, 10 \text{ km s}^{-1})$ and $(\delta\theta, \delta v) = (0.1 \text{ mas}, 2 \text{ km s}^{-1})$. However, since the parameter uncertainties scale with measurement error σ and number of monitored stars N as $\sigma/N^{1/2}$, the results can be used to describe the capabilities of an ELT with different specifications. For example, a 100 meter ELT will detect $\sim 10\times$ as many stars (§ 2.2); if the telescope has astrometric and spectroscopic errors that are smaller than those of a 30 meter telescope by a factor of five the parameter uncertainties will be $\sim 10\times$ smaller. In this section, we estimate the limits that can be placed on the parameters associated with the black hole including M_{bh} and R_0 (§ 5.1), as well as on the extended distribution of (dark) mat-




FIG. 3.— The constraints on M_{bh} and R_0 obtainable with an ELT assuming an extended matter distribution with $\gamma = 1.5$ (results are similar for $\gamma = 2$). The solid contours show the 68%, 95%, and 99.7% confidence levels assuming an astrometric limit of $\delta\theta = 0.5 \text{ mas}$ and a spectroscopic limit of $\delta v = 10 \text{ km s}^{-1}$ for the draw of 20 stars shown in Table 1. The line-dot contour shows the 99.7% confidence level for a different draw of 20 stars. The dashed contour shows the 99.7% confidence level for smaller astrometric and spectroscopic limits of $\delta\theta = 0.1 \text{ mas}$ and $\delta v = 2 \text{ km s}^{-1}$. An ELT will constrain both M_{bh} and R_0 to better than 0.1%.




FIG. 4.— The constraint on the extended matter distribution obtainable with an ELT. Shown are the 68%, 95%, and 99.7% confidence levels on the enclosed mass and slope of an extended matter distribution assuming an astrometric limit of $\delta\theta = 0.5 \text{ mas}$ and a spectroscopic limit of $\delta v = 10 \text{ km s}^{-1}$. The input models have power-law slope of $\gamma = 1.5$ and $\gamma = 2$ and an input enclosed mass of $6000 M_\odot$ within 0.01 pc. The dashed contour is the constraint at the 99.7% level for measurement errors that are a factor of five smaller.

ter near the black hole (§ 5.2). We discuss the dependence of the limits on the astrometric and spectroscopic precision of the observations. We also investigate whether relativistic




FIG. 5.— An ELT’s sensitivity to post-Newtonian effects assuming an astrometric limit of $\delta\theta = 0.5$ mas and a spectroscopic limit of $\delta v = 10 \text{ km s}^{-1}$. Shown is the uncertainty in the speed of light and the extended matter mass as obtained by including post-Newtonian corrections to the equations of motion. The Roemer delay and special relativistic effects are not included in the model in order to demonstrate that general relativistic effects of order $(v/c)^2$, including the prograde precession, are detectable with an ELT. The line styles are the same as in Figure 3.

corrections to the Keplerian motion can be detected at the GC (§ 5.3).

5.1. Measuring M_{bh} and R_0

In Figure 3 we show the constraints an ELT will place on M_{bh} and R_0 . For an astrometric limit of $\delta\theta = 0.5$ mas and a spectroscopic limit of $\delta v = 10 \text{ km s}^{-1}$ (see § 2.1) the fractional uncertainties in M_{bh} and R_0 are less than 0.1% at the 99.7% level. This is a factor of ~ 100 times better than present uncertainties. The result is robust in that simulations with distinct mock data sets of 20 stars, drawn in the fashion described in § 4.2, produce similar uncertainties in the model parameters.

For astrometric and spectroscopic limits that are a factor of five smaller the fractional uncertainties in M_{bh} and R_0 are smaller by almost a factor of five. The uncertainties in M_{bh} and R_0 scale almost linearly with the measurement errors for observations at this precision. We also verified that the uncertainties scale with N as roughly $N^{-1/2}$.

Observations with a 30 meter ELT will therefore constrain the distance to the GC to within a few parsecs and the mass of the black hole to within a few thousand solar masses. We discuss the implications of measuring R_0 to such high accuracy in § 6.

5.2. Measuring the Extended Matter Distribution

In Figure 4 we show the constraints an ELT will place on the extended matter distribution for input power-law models with $M_{\text{ext}}(r < 0.01 \text{ pc}) = 6000 M_\odot$ and $\gamma = 1.5$ or $\gamma = 2$. We chose these distributions in order to conform to the extrapolation of the observed stellar density distribution and to theoretical estimates of dark matter clustering (see § 1). We find that for an

ELT with $\delta\theta = 0.5$ mas and $\delta v = 10 \text{ km s}^{-1}$ one can detect such extended matter distributions, yielding measurements of M_{ext} and γ that are accurate to 20–30% (i.e., $\delta M_{\text{ext}} \sim 1500 M_\odot$ and $\delta\gamma \sim 0.5$). Since the amplitude of the Newtonian retrograde precession varies linearly with M_{ext} (§ 3.2), the fractional uncertainty is $\delta M_{\text{ext}}/M_{\text{ext}} \propto \delta\Delta\phi_{\text{Newt}}/\Delta\phi_{\text{Newt}} \propto M_{\text{ext}}^{-1}$, where $\delta\Delta\phi_{\text{Newt}}$ is set by the astrometric precision. Thus δM_{ext} is independent of M_{ext} so that an extended matter distribution is detectable (i.e., observations yield a lower bound) for $\delta\theta = 0.5$ mas and $\delta v = 10 \text{ km s}^{-1}$ as long as $M_{\text{ext}}(r < 0.01 \text{ pc}) \gtrsim \delta M_{\text{ext}} \sim 1500 M_\odot$. Such an ELT will therefore place interesting constraints on the extended matter at the GC.

5.3. Measuring Relativistic Effects

As discussed in § 3.3, order of magnitude estimates suggest that post-Newtonian corrections to the equations of motion, involving terms of order $(v/c)^2$, are measurable with an ELT with astrometric resolution of $\delta\theta \lesssim 0.5$ mas. In an effort to demonstrate this more quantitatively, we allow the speed of light to be a parameter in our model and examine how well we recover its value. We purposely do not include relativistic corrections to the observed motion associated with propagation effects (e.g., the Roemer time delay and other higher-order corrections) so that we can examine the detectability of $(v/c)^2$ general relativistic corrections to the orbital dynamics such as the prograde precession of the major axis position. In Figure 5 we show the constraint on c as a function of M_{enc} . Post-Newtonian effects are observable, as c is measured to $\sim 5\%$ accuracy. Since $v/c \lesssim 0.2$ for all stars in the sample (§ 3.1) the few percent constraint on c suggests that while the $(v/c)^2$ effects are measurable, the $(v/c)^3$ effects are not. The orbital precession due to black hole spin is of order $(v/c)^3$ (§ 3.4) and detecting it with an ELT with $\delta\theta = 0.5$ mas requires the favorable discovery of a star on a compact and highly eccentric orbit. Based on estimates of the signal-to-noise from a spin-induced apocenter shift (equation [15]), an astrometric precision of ~ 0.05 mas is needed to reliably detect the black hole spin.

The degeneracy between c and M_{enc} is a consequence of the degeneracy between the prograde relativistic precession and the retrograde Newtonian precession. Decreasing c increases the amount of prograde motion $\Delta\phi_{\text{pro}}$ while increasing M_{enc} increases the amount of retrograde motion $\Delta\phi_{\text{Newt}}$. The two effects compensate for one another over a range of c and M_{enc} . The degeneracy is broken at sufficiently extreme values of M_{enc} because the relativistic and Newtonian effects each induce a distinct precessional shape.

6. CONSTRAINTS ON GALACTIC STRUCTURE FROM MEASUREMENTS OF R_0

The distance to the GC, R_0 , is a fundamental parameter in models of the Milky Way structure. As Olling & Merrifield (2001) note, models of the Milky Way exhibit strong interrelations between the Galactic constants (R_0 and the local Galactic rotation speed Θ_0), the shortest-to-longest axis ratio, $q = c/a$, of the dark matter halo, and the local stellar column density Σ_* . The determination of q is of particular interest since different models of dark-matter and structure formation scenarios predict different values for q . Cold dark-matter simulations typically produce galactic halos that are triaxial (Warren et al. 1992; Jing & Suto 2002) although these tend become oblate under the influence of the dissipative infall of gas resulting in halos with $q \simeq 0.5$ (Dubinski 1994). Alternatively, hot dark-matter models predict round halos with

$q \sim 0.8$ (Peebles 1993) while some baryonic dark matter models imply $q \sim 0.2$ (Pfenniger, Combes, & Martinet 1994). As we now discuss, determining R_0 to 0.1% via monitoring of stellar orbits at the GC with an ELT enables an extremely precise measurement of q in the Milky Way.

Olling & Merrifield (2000) demonstrate that there is significant uncertainty in existing estimates of q in galaxies due to both the limited amount of data available for measuring q and the fact that different measurement techniques have yielded systematically different values. Presently, the situation is not any better for our own Galaxy, with plausible values lying in the range $0.3 \lesssim q \lesssim 1$.

The measurement of q in the Milky Way entails measuring the Galaxy's radial mass distribution and the degree to which this mass distribution is flattened. Olling & Merrifield (2000) show that the uncertainty in q in the Milky Way is almost entirely due to the large errors in the Galactic constants Θ_0 and R_0 . Indeed, Olling & Merrifield (2001) show that the fractional uncertainty in q is nearly twice the fractional uncertainty in Θ_0 . Therefore, a precision measurement of the Sun's proper motion with respect to the GC in combination with a precision measurement of R_0 tightly constrains Θ_0 and hence q . According to Salim, Gould, & Olling (2002) future astrometric surveys will be able to measure the Sun's proper motion $\mu = V/R_0$ to within several microarcseconds, corresponding to 0.1% accuracy. Here $V = \Theta_0 + V_\odot \simeq 220 \text{ km s}^{-1}$ is the sum of the rotation speed of the local standard of rest and the Sun's motion relative to it. The uncertainty in Θ_0 will be the dominant error in V ; $V_\odot$ is already known to an accuracy of 0.6 km s^{-1} from the *Hipparcos* catalogue (Dehnen & Binney 1998). Thus, the monitoring of stellar orbits at the GC with an ELT in conjunction with future astrometric survey missions will constrain the Milky Way's dark matter halo shape parameter q to a few tenths of a percent.

7. CONCLUSIONS

We have examined a variety of experiments that can be achieved through the infrared monitoring with an ELT of stars within a few thousand AU of the GC. The astrometric limit of a 30 meter ELT is conservatively 0.5 mas and possibly as high as 0.1 mas. By comparison, the astrometric limit of current observations is 1–2 mas.

The greater point-source sensitivity and spectral resolution of an ELT enables the measurement of radial velocities with errors $\lesssim 10 \text{ km s}^{-1}$. At present, of the ~ 10 stars with measured accelerated proper motions, spectral lines have been detected only in S0-2, with radial velocity uncertainties of $\sim 30 \text{ km s}^{-1}$. Measuring the radial velocities of stars breaks the degeneracy between mass and distance and thus yields a direct measurement of the distance to the GC. If the spectra of fainter stars can be obtained, the detection of deep molecular lines will improve upon the velocity estimates by an additional factor $\times 10$. The solar type stars that will be detectable with an ELT may therefore yield radial velocity uncertainties considerably smaller than 10 km s^{-1} .

A 30 meter ELT will be able to detect stars down to a K -band magnitude of $K \sim 22$, approximately four magnitudes fainter than currently possible. Due to confusion, it will be difficult to detect still fainter stars. Using measurements of the K -band luminosity function within the inner $1''$ of the GC, we estimate that such an ELT will detect the accelerated motion of ~ 100 stars with semi-major axes in the range $200 \lesssim a \lesssim 3000 \text{ AU}$. Current observations are limited to the

detection of ~ 10 stars, all with $a \gtrsim 1000 \text{ AU}$. We find that the number of stars with detectable accelerated motion scales with the aperture of an ELT as $N \simeq 100(D/30 \text{ m})^2$.

Given the observational capabilities of an ELT and the likely, albeit at low masses largely uncertain, stellar environment at the GC, we constructed a plausible sample of stellar orbits. The model includes the dynamical contribution of an extended distribution of dark matter around the black hole that is composed of stellar remnants and CDM. We find that for measurements at the precision obtainable with an ELT the uncertainty in the model parameters scale with the measurement errors σ (i.e., $\delta\theta$, δv) and the number of monitored stars N as roughly $\sigma/N^{1/2}$. Thus, while we focus on the capabilities of a diffraction limited 30 meter ELT with $\delta\theta = 0.5 \text{ mas}$ and $\delta v = 10 \text{ km s}^{-1}$, our results can be used to determine the capabilities of an ELT with different specifications. For example, a 100 meter ELT will detect $\sim 10\times$ as many stars so that if it has astrometric and spectroscopic errors that are smaller by a factor of five, the measurement accuracy in the parameters will improve by a factor of approximately ten.

We find that with a 30 meter ELT the parameters M_{bh} and R_0 will be measured to an accuracy better than 0.1%. Determining R_0 to within a few parsecs will significantly constrain models of the Galactic structure as it aids the precise measurement of the dark matter halo shape.

While current observations of stellar proper motions are compatible with Keplerian motion, a number of dynamical effects produce significant deviations, including the Newtonian retrograde precession, the relativistic prograde precession, frame dragging due to the black hole spin, and interstellar interactions involving nearby encounters. All but the frame dragging effect produce non-Keplerian motions that are detectable with a 30 meter ELT. Unfortunately, the spin of the massive black hole at the GC will probably be out of reach to kinematic studies unless an astrometric precision of $\sim 0.05 \text{ mas}$ is achieved.

The presence of an extended distribution of matter results in a Newtonian retrograde precession due to differences in the amount of mass enclosed within an orbit's pericenter and apocenter. We considered extended matter density profiles consistent with current observations of the stellar distribution at the GC. We modeled the distribution as a power-law profile normalized such that $M_{\text{ext}}(r < 0.01 \text{ pc}) = 6000M_\odot$ and with slope $\gamma = 1.5$ or 2. Standard models of dark matter clustering about a massive black hole predict similar profiles. An orbit monitoring program with a 30 meter ELT will constrain the mass and slope of such profiles to $\sim 30\%$ accuracy. Thus, monitoring orbits with an ELT provides a probe of the extended matter distribution within $\sim 10^4$ Schwarzschild radii of the massive black hole at the GC.

We also calculated the rate at which the monitored stars experience detectable deflections due to stellar gravitational scattering encounters with background compact remnants. We considered a detection threshold set by the minimum detectable change in the velocity of a monitored star. For a density cusp dominated by $\sim 10M_\odot$ black holes, ~ 30 nearby stellar encounters will be detected by a 30 meter ELT over a ten year observing baseline. This will confirm the presence of a cusp of compact remnants at the GC and enable the measurement of the remnants' masses.

The authors would like to thank M. Kamionkowski, D. Figer, K. Matthews, and E. Pfahl for comments and

E. Agol, S. Phinney, and J. Graham for helpful discussions. We also thank the referees for their helpful comments. NNW acknowledges the support of an NSF graduate fellowship

and DoE DE-FG03-92ER40701. MM was supported at Caltech by a postdoctoral fellowship from the Sherman Fairchild Foundation.

REFERENCES

- Alexander, T. & Sternberg, A. 1999, *ApJ*, 520, 137
 Backer, D. C. & Sramek, R. A. 1999, *ApJ*, 524, 805
 Baganoff, F. K. et al. 2001, *Nature*, 413, 45
 Baganoff, F. K. et al. 2003, *ApJ*, 591, 891
 Bahcall, J. N. & Wolf, R. A. 1976, *ApJ*, 209, 214
 Bahcall, J. N. & Wolf, R. A. 1977, *ApJ*, 216, 883
 Beckert, T., Duschl, W. J., Mezger, P. G., & Zylka, R. 1996, *A&A*, 307, 450
 Becklin, E. E. 1968, Ph.D. Thesis,
 Dehnen, W. & Binney, J. J. 1998, *MNRAS*, 298, 387
 Dubinski, J. 1994, *ApJ*, 431, 617
 Eckart, A., Genzel, R., Hofmann, R., Sams, B. J., & Tacconi-Garman, L. E. 1995, *ApJ*, 445, L23
 Eckart, A., Genzel, R., Ott, T., & Schödel, R. 2002, *MNRAS*, 331, 917
 Eisenhauer, F., Schödel, R., Genzel, R., Ott, T., Tecza, M., Abuter, R., Eckart, A., & Alexander, T. 2003, *ApJ*, 597, L121
 Figer, D. F. et al. 2000, *ApJ*, 533, L49
 Figer, D. F. et al. 2003, *ApJ*, 599, 1139
 Fragile, P. C. & Mathews, G. J. 2000, *ApJ*, 542, 328
 Franceschini, A., Toffolatti, L., Danese, L., & de Zotti, G. 1989, *ApJ*, 344, 35
 Gelman, A., Carlin, J. B., Stern, H. S., & Rubin, D. B. 1995, *Bayesian Data Analysis* (New York: Chapman & Hall/CRC)
 Genzel, R., Pichon, C., Eckart, A., Gerhard, O. E., & Ott, T. 2000, *MNRAS*, 317, 348
 Genzel, R. et al. 2003a, *ApJ*, 594, 812
 Genzel, R., Schödel, R., Ott, T., Eckart, A., Alexander, T., Lacombe, F., Rouan, D., & Aschenbach, B. 2003b, *Nature*, 425, 934
 Gerhard, O. 2001, *ApJ*, 546, L39
 Gezari, S., Ghez, A. M., Becklin, E. E., Larkin, J., McLean, I. S., & Morris, M. 2002, *ApJ*, 576, 790
 Ghez, A. M., Klein, B. L., Morris, M., & Becklin, E. E. 1998, *ApJ*, 509, 678
 Ghez, A. M. et al. 2003a, *ApJ*, 586, L127
 Ghez, A. M., Salim, S., Hornstein, S. D., Tanner, A., Morris, M., Becklin, E. E., & Duchêne, G. 2003b, preprint (astro-ph/0306130)
 Ghez, A. M., Wright, S. A., Matthews, K., Thompson, D., Le Mignant, D., Tanner, A., Hornstein, S. D., Morris, M., Becklin, E. E., Soifer, B. T. 2004, *ApJ*, 601, L159
 Gilks, W. R., Richardson, S., & Spiegelhalter, D. J. 1996, *Markov Chain Monte Carlo in Practice* (New York: Chapman & Hall/CRC)
 Gnedin, O. Y. & Primack, J. R. 2003, preprint (astro-ph/0308385)
 Goldwurm, A., Brion, E., Goldoni, P., Ferrando, P., Daigne, F., Decourchelle, A., Warwick, R. S., & Predehl, P. 2003, *ApJ*, 584, 751
 Gondolo, P. & Silk, J. 1999, *Physical Review Letters*, 83, 1719
 Gould, A. & Quillen, A. C. 2003, *ApJ*, 592, 935
 Hansen, B. M. S. & Milosavljević, M. 2003, *ApJ*, 593, L77
 Hornstein, S. D., Ghez, A. M., Tanner, A., Morris, M., Becklin, E. E., & Wizinowich, P. 2002, *ApJ*, 577, L9
 Ipser, J. R. & Sikivie, P. 1987, *Phys. Rev. D*, 35, 3695
 Jackson, J. M., Geis, N., Genzel, R., Harris, A. I., Madden, S., Poglitsch, A., Stacey, G. J., & Townes, C. H. 1993, *ApJ*, 402, 173
 Jaroszynski, M. 1998, *Acta Astronomica*, 48, 653
 Jing, Y. P. & Suto, Y. 2002, *ApJ*, 574, 538
 Kim, S. S. & Morris, M. 2003, *ApJ*, 597, 312
 Krabbe, A. et al. 1995, *ApJ*, 447, L95
 Landau, L. & Lifshitz, D. 1960, *Mechanics* (Oxford: Pergamon Press)
 Levin, Y. & Beloborodov, A. M. 2003, *ApJ*, 590, L33
 Loeb, A. 2003, preprint (astro-ph/0309716)
 Magorrian, J. & Tremaine, S. 1999, *MNRAS*, 309, 447
 Merritt, D., Milosavljević, M., Verde, L., & Jimenez, R. 2002, *Physical Review Letters*, 88, 191301
 Merritt, D. 2003, preprint (astro-ph/0311594)
 Metropolis, N., Rosenbluth, A. W., Rosenbluth, M. N., Teller, A. H., Teller, E. 1953, *J. Chem. Phys.*, 21, 1087
 Milosavljević, M. & Loeb, A. 2004, 604, L45
 Miralda-Escudé, J. & Gould, A. 2000, *ApJ*, 545, 847
 Morris, M. 1993, *ApJ*, 408, 496
 Munyaneza, F., Viollier, R. D. 1999, preprint (astro-ph/9910566)
 Murphy, B. W., Cohn, H. N., & Durisen, R. H. 1991, *ApJ*, 370, 60
 Murray, C. D., Dermott, S. F. 1999, *Solar System Dynamics* (Cambridge: Cambridge University Press)
 Olling, R. P. & Merrifield, M. R. 2000, *MNRAS*, 311, 361 (OM00)
 Olling, R. P. & Merrifield, M. R. 2001, *MNRAS*, 326, 164
 Peebles, P. J. E. 1972, *ApJ*, 178, 371
 Peebles, P. J. E. 1993, *Principles of Physical Cosmology* (Princeton, NJ: Princeton University Press)
 Pfahl, E. & Loeb, A. 2003, preprint (astro-ph/0309744)
 Pfenniger, D., Combes, F., & Martinet, L. 1994, *A&A*, 285, 79
 Porquet, D., Predehl, P., Aschenbach, B., Grosso, N., Goldwurm, A., Goldoni, P., Warwick, R. S., & Decourchelle, A. 2003, *A&A*, 407, L17
 Quinlan, G. D., Hernquist, L., & Sigurdsson, S. 1995, *ApJ*, 440, 554
 Reid, M. J. 1993, *ARA&A*, 31, 345
 Reid, M. J., Readhead, A. C. S., Vermeulen, R. C., & Treuhaft, R. N. 1999, *ApJ*, 524, 816
 Rogers, A. E. E. et al. 1994, *ApJ*, 434, L59
 Salim, S., Gould, A., & Olling, R. P. 2002, *ApJ*, 573, 631
 Rieke, G. H., Rieke, M. J., & Paul, A. E. 1989, *ApJ*, 336, 752
 Rubilar, G. F. & Eckart, A. 2001, *A&A*, 374, 95
 Sanders, R. H. & Lowinger, T. 1972, *AJ*, 77, 292
 Schödel, R. et al. 2002, *Nature*, 419, 694
 Schödel, R., Ott, T., Genzel, R., Eckart, A., Mouawad, N., & Alexander, T. 2003, *ApJ*, 596, 1015
 Serabyn, E., Carlstrom, J., Lay, O., Lis, D. C., Hunter, T. R., & Lacy, J. H. 1997, *ApJ*, 490, L77
 Shao, M. & Colavita, M. M. 1992, *A&A*, 262, 353
 Spitzer, L. 1987, *Dynamical evolution of globular clusters* (Princeton, NJ: Princeton University Press)
 Takeuchi, T. T., Kawabe, R., Kohno, K., Nakanishi, K., Ishii, T. T., Hirashita, H., & Yoshikawa, K. 2001, *PASP*, 113, 586
 Warren, M. S., Quinn, P. J., Salmon, J. K., & Zurek, W. H. 1992, *ApJ*, 399, 405
 Weinberg, S. 1972, *Gravitation and cosmology: Principles and applications of the general theory of relativity* (New York: Wiley)
 Ullio, P., Zhao, H., & Kamionkowski, M. 2001, *Phys. Rev. D*, 64, 43504
 Young, P. 1980, *ApJ*, 242, 1232
 Yu, Q. 2003, *MNRAS*, 339, 189
 Zhao, J., Bower, G. C., & Goss, W. M. 2001, *ApJ*, 547, L29